


=>

Uploading C:\Program Files\Stnexp\Queries\10551564-amended.str



chain nodes :

11 13 14 15 16 17

ring nodes :

1 2 3 4 5 6 7 8 9 18 19 20 21 22 23 24 25 26 27 28 29

chain bonds :

7-11 8-18 9-27 11-13 13-14 13-15 15-16 16-17

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-9 7-8 8-9 18-19 18-23 19-20 20-21 21-22  
22-23 24-25 24-29 25-26 26-27 27-28 28-29

exact/norm bonds :

5-6 5-7 6-9 7-8 7-11 8-9 8-18 9-27 11-13 13-14 13-15 15-16 16-17

exact bonds :

24-25 24-29 25-26 26-27 27-28 28-29

normalized bonds :

1-2 1-6 2-3 3-4 4-5 18-19 18-23 19-20 20-21 21-22 22-23

isolated ring systems :

containing 1 : 18 : 24 :

Connectivity :

11:2 E exact RC ring/chain 16:2 E exact RC ring/chain

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 11:CLASS  
13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom  
22:Atom 23:Atom  
24:Atom 25:Atom 26:Atom 27:Atom 28:Atom 29:Atom

L1 STRUCTURE UPLOADED

=> d his

FILE 'REGISTRY' ENTERED AT 15:28:10 ON 18 AUG 2008

L1 STRUCTURE UPLOADED


L3 48 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 15:29:20 ON 18 AUG 2008

L4 8 S L3

FILE 'REGISTRY' ENTERED AT 15:29:32 ON 18 AUG 2008

=> d l1



=> fil caplus

=> d 14 tot bib abs hitstr

√<sub>L4</sub> ANSWER 1 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN  
 AN 2007:923552 CAPLUS Full-text  
 DN 147:448607  
 TI Development of carboxylic acid replacements in indole-N-acetamide  
 inhibitors of hepatitis C virus NS5B polymerase  
 AU Stansfield, Ian; Pompei, Marco; Conte, Immacolata; Ercolani, Caterina;  
 Migliaccio, Giovanni; Jairaj, Mark; Giuliano, Claudio; Rowley, Michael;  
 Narjes, Frank  
 CS IRBM (Merck Research Laboratories Rome), Rome, 00040, Italy  
 SO Bioorganic & Medicinal Chemistry Letters √(2007), 17(18), 5143-5149  
 CODEN: BMCLE8; ISSN: 0960-894X  
 PB Elsevier Ltd.  
 DT Journal  
 LA English  
 OS CASREACT 147:448607  
 GI



AB Allosteric inhibition of the hepatitis C virus (HCV) NS5B RNA-dependent RNA polymerase enzyme has recently emerged as a viable strategy toward blocking replication of viral RNA in cell-based systems. We report here 2 series of indole-N-acetamides, bearing physicochem. diverse carboxylic acid replacements, which show potent affinity for the NS5B enzyme with reduced potential for formation of glucuronide conjugates. E.g., indole-N-acetamide I was prepared in several steps from Me 2-bromo-3-cyclohexyl-5-indolecarboxylate. Preliminary optimization of these series furnished compds. that are potent in the blockade of subgenomic HCV RNA replication in HUH-7 cells.


IT 774213-31-9P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of indole-N-acetamides as inhibitors of hepatitis C virus NS5B polymerase)

RN 774213-31-9 CAPLUS

CN 1H-Indole-1-acetamide, 3-cyclohexyl-6-(2,5-dihydro-5-oxo-1,2,4-oxadiazol-3-yl)-N-methyl-N-[(1-methyl-3-piperidinyl)methyl]-2-phenyl- (CA INDEX NAME)



√L4 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2007:324208 CAPLUS Full-text

DN 147:497550

TI Attenuating pregnane X receptor (PXR) activation: a molecular modelling approach

AU Gao, Y.-D.; Olson, S. H.; Balkovec, J. M.; Zhu, Y.; Royo, I.; Yabut, J.; Evers, R.; Tan, E. Y.; Tang, W.; Hartley, D. P.; Mosley, R. T.

CS CIBE, Department of Molecular Systems, Merck Research Laboratories,  
Madrid, Spain

SO Xenobiotica ✓ (2007), 37(2), 124-138

CODEN: XENOBH; ISSN: 0049-8254

PB Informa Healthcare

DT Journal

LA English

AB Recent studies have demonstrated that the pregnane X receptor (PXR) is a key regulator of cytochromes P 450 3A (e.g. CYP3A4 in human) gene expression. As a result, activation of PXR may lead to CYP3A4 protein over-expression. Because induction of CYP3A4 could result in clin. important drug-drug interactions, there has been a great interest in reducing the possibility of PXR activation by drug candidates in drug-discovery programs. In order to provide structural insight for attenuating drug candidate-mediated PXR activation, we used a docking approach to study the structure-activity relationship for PXR activators. Based on our docking models, it is proposed that introducing polar groups to the end of an activator should reduce its human PXR (hPXR) activity via destabilizing interactions in the hydrophobic areas of the PXR ligand-binding pocket. A number of analogs that incorporate these structural features then were designed and synthesized, and they exhibited significantly lower hPXR activation in a transactivation assay and decreased CYP3A4 induction in a human hepatocytes-based assay. In addition, an example in which attenuating hPXR activation was achieved by sterically destabilizing the helices 11 and 12 of the receptor is presented.


IT 774210-59-2 861966-03-2

RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(mol. modeling of pregnane X receptor (PXR) regulation)


RN 774210-59-2 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)




RN 861966-03-2 CAPLUS

CN 1H-Indole-6-carboxylic acid, 2-(4-chlorophenyl)-3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]- (CA INDEX NAME)




✓L4 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN  
 AN 2006:834581 CAPLUS Full-text  
 DN 145:410005  
 TI Parallel Screening: A Novel Concept in Pharmacophore Modeling and Virtual Screening  
 AU Steindl, Theodora M.; Schuster, Daniela; Laggner, Christian; Langer, Thierry  
 CS Institute of Pharmacy, Computer Aided Molecular Design Group, University of Innsbruck, Innrain, Austria  
 SO Journal of Chemical Information and Modeling ✓ (2006), 46(5), 2146-2157  
 CODEN: JCISD8; ISSN: 1549-9596  
 PB American Chemical Society  
 DT Journal  
 LA English  
 AB Parallel screening comprises a novel in silico method to predict the potential biol. activities of a compound by screening it with a multitude of pharmacophore models. Our aim is to provide a fast, large-scale system that allows for virtual activity profiling. In this proof of principle study, carried out with the software tools LigandScout and Catalyst, the authors present a model work for the application of parallel pharmacophore-based virtual screening on a set of 50 structure-based pharmacophore models built for various viral targets and 100 antiviral compds. The latter were screened against all pharmacophore models in order to determine if their biol. targets could be correctly predicted via an enrichment of corresponding pharmacophores matching these ligands. The results demonstrate that the desired enrichment, i.e., successful virtual activity profiling, was achieved for approx. 90% of all input mols. The authors discuss descriptors for output validation, as well as various aspects influencing the anal. of the obtained activity profiles, and the effect of the utilized search modus for screening.  
 IT 912462-96-5  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (parallel screening, a concept in pharmacophore modeling and virtual screening)  
 RN 912462-96-5 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(3R)-1-methyl-3-piperidinyl]methyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)

Absolute stereochemistry.



✓L4 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN  
 AN 2006:272517 CAPLUS Full-text  
 DN 144:311906  
 TI Preparation of indoleacetamides as antivirals for treatment of hepatitis C infection.  
 IN Colarusso, Stefania; Conte, Immacolata; Habermann, Joerg; Narjes, Frank; Ponzi, Simona  
 PA Istituto di Ricerche di Biologia Molecolare P. Angeletti S.p.A., Italy  
 SO PCT Int. Appl., 58 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006029912	A1	20060323	WO 2005-EP52631	20050608
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	AU 2005284248	A1	20060323	AU 2005-284248	20050608
	CA 2568832	A1	20060323	CA 2005-2568832	20050608
	EP 1758857	A1	20070307	EP 2005-811114	20050608
	R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, LV				
	CN 1964944	A	20070516	CN 2005-80019084	20050608
	JP 2008501767	T	20080124	JP 2007-526425	20050608
	IN 2006DN07856	A	20070817	IN 2006-DN7856	20061226
PRAI	GB 2004-13087	A	20040611		
	WO 2005-EP52631	W	✓20050608		
OS	MARPAT 144:311906				
GI					



AB Title compds. [I; E = H, alkyl, alkenyl, alkynyl, cycloalkyl, (substituted) (hetero)aryl; A = (substituted) alkyl, alkenyl, nonarom. ring; R1, R2 = H, alkyl, alkenyl, alkynyl, alkoxy, cycloalkylalkyl, etc.; R3 = H, alkyl, alkenyl; R4 = H, alkyl; X = DBCR5R6; R5, R6 = H, halo, alkyl, alkenyl, alkoxy; B = (substituted) aryl, heteroaryl, etc.; D = bond, alkylene, alkenylene, alkynylene, (substituted) aryl, heteroaryl], were prepared Thus, title compound (2E)-3-[4-[[[1-[[[3-cyclohexyl-1-[2- (dimethylamino)-2-oxoethyl]-2-phenyl-1H-indol-6-yl]carbonyl]amino]cyclopentyl]carbonyl]amino]phenyl]acrylic acid was prepared in 8 steps from Me indole-6-carboxylate, cyclohexanone, phenylboronic acid, 2-chloro-N,N-dimethylacetamide, 1-[[[(benzyloxy)carbonyl]amino]cyclopentanecarboxylic acid, and Et cinnamate. I generally showed IC50's of <1  $\mu$ M for inhibition of HCV RNA dependent RNA polymerase (NS5B).

IT 879498-47-2P 879498-61-0P 879498-65-4P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(claimed compound; preparation of indoleacetamides as antivirals for treatment of hepatitis C infection)

RN 879498-47-2 CAPLUS

CN 2-Propenoic acid, 3-[4-[[[1-[[[3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-1H-indol-6-yl]carbonyl]amino]cyclopentyl]carbonyl]amino]phenyl]-, (2E)- (CA INDEX NAME)

Double bond geometry as shown.




RN 879498-61-0 CAPLUS

CN 2-Propenoic acid, 3-[4-[[[1-[[[3-cyclohexyl-1-[2-oxo-2-[[[5-oxo-2-pyrrolidinyl)methyl]amino]ethyl]-2-phenyl-1H-indol-6-yl]carbonyl]amino]cyclopentyl]carbonyl]amino]phenyl]-, (2E)- (CA INDEX NAME)

NAME)


Double bond geometry as shown.



RN 879498-65-4 CAPLUS

CN 2-Propenoic acid, 3-[4-[[[1-[[[3-cyclohexyl-1-[2-[(1,4-dioxan-2-ylmethyl)amino]-2-oxoethyl]-2-phenyl-1H-indol-6-yl]carbonyl]amino]cyclopentyl]carbonyl]amino]phenyl]-, (2E)- (CA INDEX NAME)

Double bond geometry as shown.



√<sub>L4</sub> ANSWER 5 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2005:760643 CAPLUS [Full-text](#)


DN 143:341687

TI Interdomain Communication in Hepatitis C Virus Polymerase Abolished by Small Molecule Inhibitors Bound to a Novel Allosteric Site

AU Di Marco, Stefania; Volpari, Cinzia; Tomei, Licia; Altamura, Sergio; Harper, Steven; Narjes, Frank; Koch, Uwe; Rowley, Michael; De Francesco,




Raffaele; Migliaccio, Giovanni; Carfi, Andrea  
 CS Istituto di Ricerche di Biologia Molecolare P. Angeletti, Pomezia (Rome),  
 00040, Italy  
 SO Journal of Biological Chemistry  $\sqrt{\quad}$  (2005), 280(33), 29765-29770  
 CODEN: JBCHA3; ISSN: 0021-9258  
 PB American Society for Biochemistry and Molecular Biology  
 DT Journal  
 LA English  
 AB The hepatitis C virus (HCV) NS5B RNA-dependent RNA polymerase is required for  
 replication of the viral genome and is a key target for therapeutic  
 intervention against HCV. We have determined the crystal structures of the  
 HCV polymerase complexed with two indole-based allosteric inhibitors at 2.3-  
 and 2.4-Å resolution. The structures show that these inhibitors bind to a site  
 on the surface of the thumb domain. A cyclohexyl and Ph ring substituents,  
 bridged by an indole moiety, fill two closely spaced pockets, whereas a  
 carboxylate substituent forms a salt bridge with an exposed arginine side  
 chain. Interestingly, in the apoenzyme, the inhibitor binding site is  
 occupied by a small  $\alpha$ -helix at the tip of the N-terminal loop that connects  
 the fingers and thumb domains. Thus, these mols. inhibit the enzyme by  
 preventing formation of intramol. contacts between these two domains and  
 consequently precluding their coordinated movements during RNA synthesis. Our  
 structures identify a novel mechanism by which a new class of allosteric  
 inhibitors inhibits the HCV polymerase and open the way to the development of  
 novel antiviral agents against this clin. relevant human pathogen.  
 IT 774210-59-2D, complexes with NS5B  
 RL: PRP (Properties)  
 (crystal structures reveal interdomain communication in HCV NS5B  
 polymerase is disrupted by indole-based inhibitors bound to novel  
 allosteric site)  
 RN 774210-59-2 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-3-  
 piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)



$\sqrt{\quad}$ L4 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN  
 AN 2005:500671 CAPLUS Full-text  
 DN 143:211802  
 TI Potent inhibitors of subgenomic hepatitis C virus RNA replication through  
 optimization of indole-N-acetamide allosteric inhibitors of the viral NS5B  
 polymerase  
 AU Harper, Steven; Avolio, Salvatore; Pacini, Barbara; Di Filippo, Marcello;  
 Altamura, Sergio; Tomei, Licia; Paonessa, Giacomo; Di Marco, Stefania;  
 Carfi, Andrea; Giuliano, Claudio; Padron, Julio; Bonelli, Fabio;  
 Migliaccio, Giovanni; De Francesco, Raffaele; Laufer, Ralph; Rowley,  
 Michael; Narjes, Frank  
 CS IRBM Merck Research Laboratories, Rome, 00040, Italy

SO Journal of Medicinal Chemistry ✓ (2005), 48(14), 4547-4557  
 CODEN: JMCMAR; ISSN: 0022-2623  
 PB American Chemical Society  
 DT Journal  
 LA English  
 OS CASREACT 143:211802  
 GI



AB Infections caused by hepatitis C virus (HCV) are a significant world health problem for which novel therapies are in urgent demand. Compds. that block replication of subgenomic HCV RNA in liver cells are of interest because of their demonstrated antiviral effect in the clinic. In followup to a recent report that indole-N-acetamides were potent allosteric inhibitors of the HCV NS5B polymerase enzyme, the optimization as cell-based inhibitors are described. The crystal structure of I bound to NS5B was a guide in the design of a two-dimensional compound array that highlighted that formally zwitterionic inhibitors have strong intracellular potency and that pregnane X receptor (PXR) activation (an undesired off-target activity) was linked to a structural feature of the inhibitor. Optimized analogs devoid of PXR activation (EC<sub>50</sub> = 127 nM) retain strong cell-based efficacy under high serum conditions and showed acceptable pharmacokinetics parameters in rat and dog.

IT 774210-60-5P 861966-04-3P 861966-27-0P  
 861966-47-4P

RL: CPN (Combinatorial preparation); PAC (Pharmacological activity); PRP (Properties); BIOL (Biological study); CMBI (Combinatorial study); PREP (Preparation)  
 (preparation, subgenomic hepatitis C virus and viral NS5B polymerase inhibitory activity, and structure-activity relationship of indole-N-acetamide derivs. using either combinatorial chemical or solution-phase chemical)


RN 774210-60-5 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 774210-59-2


CMF C31 H39 N3 O3



CM 2

CRN 76-05-1

CMF C2 H F3 O2




RN 861966-04-3 CAPLUS

CN 1H-Indole-6-carboxylic acid, 2-(4-chlorophenyl)-3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 861966-03-2


CMF C31 H38 Cl N3 O3



CM 2

CRN 76-05-1


CMF C2 H F3 O2



RN 861966-27-0 CAPLUS  
 CN 1H-Indole-5-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)


CM 1

CRN 774212-42-9  
 CMF C31 H39 N3 O3



CM 2



CRN 76-05-1  
 CMF C2 H F3 O2



RN 861966-47-4 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-2-(3-fluorophenyl)-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1


CRN 774211-97-1  
 CMF C31 H38 F N3 O3



CM 2

CRN 76-05-1

CMF C2 H F3 O2



√L4 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN  
 AN 2005:141029 CAPLUS Full-text  
 DN 142:240430  
 TI Preparation of heterocyclic compounds as hepatitis C virus polymerase inhibitors  
 IN Oka, Takahiro; Yata, Shinji; Ikegashira, Kazutaka; Noji, Satoru; Akaki, Tatsuo; Hirashima, Shintaro; Niwa, Yasushi; Ando, Izuru; Sato, Toshihiro  
 PA Japan Tobacco Inc., Japan  
 SO PCT Int. Appl., 467 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	√DATE
PI	WO 2005014543	A1	20050217	WO 2004-JP11640	20040806
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRAI JP 2003-288296 A 20030806  
 JP 2003-288298 A 20030806

√L4 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2004:857606 CAPLUS Full-text

DN 141:350034

TI Preparation of indole acetamides as inhibitors of the hepatitis c virus  
 NS5B polymerase

IN Avolio, Salvatore; Di Filippo, Marcello; Harper, Steven; Narjes, Frank;  
 Pacini, Barbara; Pompei, Marco; Rowley, Michael; Stansfield, Ian

PA Istituto Di Ricerche Di Biologia Molecolare P Angeletti Spa, Italy

SO PCT Int. Appl., 126 pp.


CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004087714	A1	20041014	WO 2004-GB1437	20040402
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2004226144	A1	20041014	AU 2004-226144	20040402
	CA 2520896	A1	20041014	CA 2004-2520896	20040402
	EP 1613634	A1	20060111	EP 2004-725422	20040402
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	JP 2007516158	T	20070621	JP 2006-506078	20040402
	IN 2005DN04494	A	20070824	IN 2005-DN4494	20051004
	US 20070167447	A1	20070719	√US 2006-551564	20060605
PRAI	GB 2003-7891	A	20030404		
	WO 2004-GB1437	W	20040402		
OS	MARPAT 141:350034				
GI					



I



II

AB Title compds. represented by the formula I [wherein Ar1 = (un)substituted heteroaryl; A1 = (un)substituted alkyl, alkenyl, non-aromatic (bi)cyclic ring; R1, R2 = independently H, alkyl, alkenyl, alkynyl, etc.; n = 1-4; X1-X4 = N or (un)substituted carbon; and pharmaceutically acceptable salts thereof] were prepared as inhibitors of the hepatitis c virus (HCV) NS5B polymerase. For example, II was given in a multi-step synthesis starting from the reaction of Me 1H-indole-6-carboxylate with 3-bromocyclohex-1-ene. I were tested for inhibitory activity against the HCV RNA dependent RNA polymerase (NS5B) in an enzyme inhibition assay with IC50 below 5µM in the enzyme assay and EC50 below 20 pM in the cell based assay. Thus, I and their pharmaceutical compns. are useful as inhibitors of the hepatitis c virus NS5B polymerase for the prevention and treatment of hepatitis C infections.


IT 774210-51-4P 774210-57-0P 774210-60-5P  
774210-63-8P 774210-66-1P 774210-69-4P  
774210-75-2P 774211-22-2P 774211-44-8P  
774211-50-6P 774211-97-1P 774212-18-9P  
774212-22-5P 774212-42-9P 774213-31-9P  
774213-35-3P 774213-72-8P 774214-26-5P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of indole acetamides as inhibitors of hepatitis c virus NS5B polymerase)

RN 774210-51-4 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[[[(1-methyl-3-pyrrolidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-, hydrochloride (1:1) (CA INDEX NAME)




● HCl

RN 774210-57-0 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[[1-(3-methyl-1H-1,2,4-triazol-5-yl)ethyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)


CM 1

CRN 774210-56-9  
 CMF C28 H31 N5 O3



CM 2


CRN 76-05-1  
 CMF C2 H F3 O2



RN 774210-60-5 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 774210-59-2  
 CMF C31 H39 N3 O3






CM 2

CRN 76-05-1

CMF C2 H F3 O2




RN 774210-63-8 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[[ (1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)

CM 1

CRN 774210-62-7


CMF C30 H37 N3 O3



CM 2

CRN 76-05-1

CMF C2 H F3 O2




RN 774210-66-1 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-2-piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)


CM 1

CRN 774210-65-0  
 CMF C31 H39 N3 O3



CM 2


CRN 76-05-1  
 CMF C2 H F3 O2



RN 774210-69-4 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(5-methyl-1H-imidazol-2-yl)methyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)


CM 1

CRN 774210-68-3  
 CMF C29 H32 N4 O3




CM 2


CRN 76-05-1  
 CMF C2 H F3 O2




RN 774210-75-2 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[[2-(1-methyl-3-pyrrolidiny)ethyl]amino]-2-oxoethyl]-2-phenyl-, 2,2,2-trifluoroacetate (1:1) (CA INDEX NAME)  
 CM 1  
 CRN 774210-74-1  
 CMF C30 H37 N3 O3




CM 2  
 CRN 76-05-1  
 CMF C2 H F3 O2




RN 774211-22-2 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[1-(2-thiazolyl)ethyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)




RN 774211-44-8 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[[2-(4-methyl-1-piperazinyl)ethyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)





RN 774211-50-6 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[[2-(4-morpholinyl)ethyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)



RN 774211-97-1 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-2-(3-fluorophenyl)-1-[2-[methyl[(1-methyl-3-piperidinyl)methyl]amino]-2-oxoethyl]- (CA INDEX NAME)




RN 774212-18-9 CAPLUS  
 CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-2-[4-[2-(dimethylamino)-2-oxoethoxy]phenyl]-1-[2-[methyl(2-pyrazinylmethyl)amino]-2-oxoethyl]- (CA INDEX NAME)




RN 774212-22-5 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-2-(3-fluorophenyl)-1-[2-oxo-2-  
[[ (tetrahydro-1,1-dioxido-3-thienyl)methyl]amino]ethyl]- (CA INDEX NAME)




RN 774212-42-9 CAPLUS

CN 1H-Indole-5-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[(1-methyl-3-  
piperidinyl)methyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)




RN 774213-31-9 CAPLUS

CN 1H-Indole-1-acetamide, 3-cyclohexyl-6-(2,5-dihydro-5-oxo-1,2,4-oxadiazol-3-  
yl)-N-methyl-N-[(1-methyl-3-piperidinyl)methyl]-2-phenyl- (CA INDEX NAME)




RN 774213-35-3 CAPLUS

CN 1H-Indole-1-acetamide, 3-cyclohexyl-6-(2,5-dihydro-5-oxo-1,2,4-oxadiazol-3-yl)-N-[(1-methyl-3-pyrrolidinyl)methyl]-2-phenyl- (CA INDEX NAME)




RN 774213-72-8 CAPLUS

CN 1H-Indole-1-acetamide, 3-cyclohexyl-N-methyl-N-[(1-methyl-3-piperidinyl)methyl]-2-phenyl-6-[[[(phenylmethyl)sulfonyl]amino]carbonyl]- (CA INDEX NAME)



RN 774214-26-5 CAPLUS

CN 1H-Indole-6-carboxylic acid, 3-cyclohexyl-1-[2-[methyl[1-(4-pyridinyl)ethyl]amino]-2-oxoethyl]-2-phenyl- (CA INDEX NAME)



=> log hold

SESSION WILL BE HELD FOR 120 MINUTES  
STN INTERNATIONAL SESSION SUSPENDED AT 15:30:00 ON 18 AUG 2008